Superinductor-based Quantum Technologies with Ultrastrong Couplings (SiUCs)

Pol Forn-Díaz, High Energy Physics Institute (IFAE), Bellaterra (Barcelona), Spain Ioan Pop, Karlsruhe Institute of Technology, Karlsruhe, Germany Milena Grifoni, Regensburg University – Institute for Theoretical Physics, Germany Miroslav Grajcar, Institute of Physics, Slovak Academy of Sciences, Bratislava, Slovakia Nicolas Roch, Néel Institute, Centre National de la Recherche Scientifique, Grenoble, France Elisabetta Paladino, Consiglio Nationale delle Richerche, Catania, Italy

Introduction

Superconducting qubits have become one of the most promising platforms on which to build the entire span of quantum technologies. The success of this technology lies in the mastering of the interaction between light and matter, framed in field named circuit quantum electrodynamics (circuit QED), where microwave photons play the role of light, while qubits effectively behave as artificially man-made atoms.

The light-matter models used by the community of superconducting quantum circuits are inherited from atomic physics, where interaction strengths are very small and the rotating-wave approximation (RWA) is fully justified. This unavoidably represents a limitation when the whole physical model is required to accurately explain the results of experiments. The general light-matter interaction model is the quantum Rabi model (QRM):

$$\hat{\mathcal{H}}_{\text{QRM}} = \frac{\hbar\omega_q}{2}\hat{\sigma_z} + \hbar\omega_r \left(\hat{a}^{\dagger}\hat{a} + \frac{1}{2}\right) + \hbar g\hat{\sigma_x}(\hat{a} + \hat{a}^{\dagger})$$

containing much richer physics, but a very large interaction strength -also known as ultrastrong coupling (USC)- is required. Very few experiments have so far attained such levels of lightmatter interaction [Forn-Diaz2010, Niemczyk2010, Yoshihara2017]. Despite this, the interest in USC physics has grown worldwide in the last decade.

Goals of project SiUCs

SiUCs aims to answer the following questions:

1. Can non-RWA effects be completely disregarded in conventional experiments of superconducting circuits exploring quantum computation and quantum simulation applications?*

2. Are there novel physics beyond the RWA that would significantly contribute to the advancement of quantum technologies with superconducting circuits?

Project goals:

- 1. Deployment of **superinductorbased** technology to achieve USCs.
- 2. Engineer novel qubits with USC physics.
- 3. Study of **applications** of USC physics in quantum technologies.
- 4. Develop applications for superinductors such as **photodetectors**.

* Recent results in the readout of transmon qubits using resonators clearly point in this direction [Sank2016] in the context of quantum computation and qubit state readout.

Superinductor enabling technology

A superinductor exhibits **an impedance exceeding** the quantum of impedance, Two strategies exist to build superinductors:



Novel qubits for USC physics

Key point in SiUCs: Employ superinductors to obtain USCs both in open (a) and closed (b) systems.

Circuit implementations for qubit-waveguide (a) and qubit-resonator (b) systems in the USC regime using superinductors.

Theoretical models will be developed to define new experimental platforms for USC physics for dynamical control and switching.





Applications of superinductor-based circuits



SKIT

HAE

The SiUCs consortium

-6 partners: 4 experiment/2 theory.
-33% Women Pis.
-50% Young researchers.
-1 widening country.

- -Execution times: 04/20-03/23.
- -Total budget: 1.29B€, 70% experiment.

[Forn-Diaz2010] P. Forn-Díaz, et al., PRL 105, 237001 (2010)
[Niemczyk2010] T. Niemczyk et al., Nature Physics 6, 772 (2010)
[Yoshihara2017] F. Yoshihara et al., Nature Physics 13, 44 (2017)
[Sank2016] D. Sank et al., PRL 117, 190503 (2016)
[Krupko2018] Y. Krupko et al., PRB (2018)
[Maleeva2018] N. Maleeva et al., Nature Comm. 9, 3889 (2018)
[De Bernardis2018] De Bernardis et al., PARA 97, 043820 (2018)
[Znyman2015] I. Znyman, S. Florens, PRB 92, 085131 (2015).

